arXiv:1605.02545v1 [physics.class-ph] 26 Apr 2016

A note on "Measuring propagation speed of Coulomb fields" by R. de Sangro, G. Finocchiaro, P. Patteri, M. Piccolo, G. Pizzella

D. M. Gitman[1,2,3],* A. E. Shabad[1,2],† and A.A. Shishmarev [2,3]‡
[1]*P. N. Lebedev Physics Institute, Leninsky Prospekt 53, Moscow 117924, Russia*
[2]*Tomsk State University, Lenin Prospekt 36, Tomsk 634050, Russia and*
[3]*Instituto de Física, Universidade de São Paulo, CEP 05508-090, São Paulo, S. P., Brazil*

In connection with the discussion and the measurements fulfilled in Ref. [1], the full identity is demonstrated between the Feynman formula for the field of a moving charge and the Liénard-Wiechert potentials.

Keywords: Electrodynamics, moving charge

In Ref. [1] measurements were performed to decide between two approaches to the field of a moving charge: one based on the Liénard-Wiechert potentials and the other on the Feynman interpretation. The aim of the present note is to demonstrate that although apparently different physical ideas are layed into the Feynman formula [4], it is as a matter of fact mathematically identical to that found in standard text books, e. g. [2, 3], for the Liénard-Wiechert potentials both for accelerated motion of the charge and its motion with a constant speed. We believe that this observation should be taken into account and prove to be useful in the general discussion on the matter [5].

The Feynmann formula for the electric field $\mathbf{E}$ of a moving accelerated charge q is [4]

$$\mathbf{E} = \frac{q}{4\pi\epsilon_0}\left[\frac{\mathbf{e}_{r'}}{r'^2} + \frac{r'}{c}\frac{d}{dt}\left(\frac{\mathbf{e}_{r'}}{r'^2}\right) + \frac{1}{c^2}\frac{d^2}{dt^2}\mathbf{e}_{r'}\right], \tag{1}$$

We set $c = 1$ for the sake of simplicity and keep to the notations in [4]. In (1) the function

$$r' = \sqrt{x'_1 + x'^2_2 + x'^2_3} \tag{2}$$

is the modulus of vector $\mathbf{x}'$, directed from the position $\tilde{\mathbf{x}}$ of the moving charge to the observation point $\mathbf{x}$:

$$\mathbf{x}' = \mathbf{x} - \tilde{\mathbf{x}}, \tag{3}$$

and $\mathbf{e}_{r'}$ is the unit vector in the direction of $\mathbf{x}'$. The charge trajectory is given as $\tilde{\mathbf{x}} = \tilde{\mathbf{x}}(t')$, where t' is the time coordinate of the charge. Therefore, with the location of the observation point fixed, $\mathbf{x} = \mathbf{const}$., we see that $\mathbf{x}'$, as well as r', is a function solely of t'. Once the influence of the charge propagates exactly with the speed of light $c = 1$, the relation

$$r'(t') = t - t' \tag{4}$$

holds, where t is the time of observation. With Eqs. (3) and (4) the length (2) is just the distance between the position of the charge and the observation point at the moment of emission.

Relation (4) defines t as a function of t'.Then, according to the rule of differentiation of an inverse function, one has for any function $a(t')$

$$\frac{da\,(t')}{dt} = \frac{da\,(t')}{dt'}\frac{dt'}{dt} = \frac{da\,(t')}{dt'}\left(\frac{dt}{dt'}\right)^{-1},$$

*Electronic address: gitman@dfn.if.usp.br
†Electronic address: shabad@lpi.ru
‡Electronic address: a.a.shishmarev@mail.ru

where $\frac{dt}{dt'}$ follows from (4) and (3) to be

$$\frac{dt}{dt'} = 1 + \frac{dr'(t')}{dt'} = 1 + \frac{d|\mathbf{x} - \tilde{\mathbf{x}}(t')|}{dt'} = 1 - \frac{(x_i - \tilde{x}_i(t'))}{r'}\frac{d\tilde{x}_i(t')}{dt'} = 1 - \frac{(\mathbf{v}\cdot\mathbf{x}')}{r'} = 1 - (\mathbf{v}\cdot\mathbf{e}_{r'}).$$

We have used here that $\frac{d\tilde{\mathbf{x}}(t')}{dt'} = \mathbf{v}(t')$ is the instantaneous speed of the charge.

Referring to the designation $(\mathbf{e}_{r'}\bar{v}) = \kappa$ used for brevity we can now rewrite Eq. (1) as

$$\mathbf{E} = \frac{q}{4\pi\epsilon_0}\left[\frac{\mathbf{e}_{r'}}{r'^2} + \frac{r'}{(1-\kappa)}\frac{d}{dt'}\left(\frac{\mathbf{e}_{r'}}{r'^2}\right) + \frac{1}{(1-\kappa)}\frac{d}{dt'}\left(\frac{1}{(1-\kappa)}\frac{d\mathbf{e}_{r'}}{dt'}\right)\right]. \tag{5}$$

Taking into account that

$$\frac{d\mathbf{e}_{r'}}{dt'} = \frac{d}{dt'}\left(\frac{\mathbf{x}'}{r'}\right) = -\frac{\mathbf{v}}{r'} + \frac{\kappa\mathbf{x}'}{r'^2} = \frac{\kappa\mathbf{e}_{r'} - \mathbf{v}}{r'} \tag{6}$$

we calculate the second term in (5):

$$\frac{r'}{(1-\kappa)}\frac{d}{dt'}\left(\frac{\mathbf{e}_{r'}}{r'^2}\right) = \frac{3\kappa\mathbf{e}_{r'} - \mathbf{v}}{(1-\kappa)\,r'^2}. \tag{7}$$

The third term in (5) is

$$\frac{1}{(1-\kappa)}\frac{d}{dt'}\left(\frac{1}{(1-\kappa)}\frac{d\mathbf{e}_{r'}}{dt'}\right) = \frac{1}{(1-\kappa)}\frac{d}{dt'}\left(\frac{\kappa\mathbf{e}_{r'} - \mathbf{v}}{(1-\kappa)\,r'}\right). \tag{8}$$

Let us calculate the derivative of $(1-\kappa)^{-1}$:

$$\frac{d\,(1-\kappa)^{-1}}{dt'} = (1-\kappa)^{-2}\left(\frac{d\mathbf{e}_{r'}}{dt'}\cdot\mathbf{v} + \mathbf{e}_{r'}\cdot\frac{d\mathbf{v}}{dt'}\right) = (1-\kappa)^{-2}\left[\frac{\kappa^2 - v^2}{r'} + (\mathbf{e}_{r'}\cdot\mathbf{v})\right], \tag{9}$$

where $\dot{\mathbf{v}}$ is the acceleration of the charge. Then the third term in (5) becomes

$$\frac{1}{(1-\kappa)}\frac{d}{dt'}\left(\frac{1}{(1-\kappa)}\frac{d\mathbf{e}_{r'}}{dt'}\right) = \frac{\left(\kappa^2 - v^2\right)(\kappa\mathbf{e}_{r'} - \mathbf{v}) + (1-\kappa)\left[2\kappa\,(\kappa\mathbf{e}_{r'} - \mathbf{v}) + \mathbf{e}_{r'}\left(\kappa^2 - v^2\right)\right]}{(1-\kappa)^3\,r'^2} +$$
$$+ \frac{(\kappa\mathbf{e}_{r'} - \mathbf{v})\,(\mathbf{e}_{r'}\cdot\mathbf{v}) + \left[\mathbf{e}_{r'}\left(\mathbf{e}_{r'}\cdot\dot{\mathbf{v}}\right) - \dot{\mathbf{v}}\right](1-\kappa)}{(1-\kappa)^3\,r'}. \tag{10}$$

Finally, substituting (7) and (10) in (5) and separating the factor $(1-\kappa)^3\,r'^2$, we get

$$\begin{aligned}\mathbf{E} = \frac{q}{4\pi\epsilon_0\,(1-\kappa)^3\,r'^2}&\left[\mathbf{e}_{r'}\,(1-\kappa)^3 + (3\kappa\mathbf{e}_{r'} - \mathbf{v})\,(1-\kappa)^2 + \right.\\ &+ \left(\kappa^2 - v^2\right)(\kappa\mathbf{e}_{r'} - \mathbf{v}) + 2\kappa\,(\kappa\mathbf{e}_{r'} - \mathbf{v})\,(1-\kappa) + \mathbf{e}_{r'}\left(\kappa^2 - v^2\right)(1-\kappa) + \\ &+ \left.r'\,(\kappa\mathbf{e}_{r'} - \mathbf{v})\left(\mathbf{e}_{r'}\cdot\dot{\mathbf{v}}\right) + r'\left[\mathbf{e}_{r'}\left(\mathbf{e}_{r'}\cdot\dot{\mathbf{v}}\right) - \dot{\mathbf{v}}\right](1-\kappa)\right].\end{aligned} \tag{11}$$

It is easy to show that this is reduced to

$$\begin{aligned}\mathbf{E} = \frac{q}{4\pi\epsilon_0\,(1-\kappa)^3\,r'^2}&\left[\mathbf{e}_{r'}\left(1-v^2\right) - \mathbf{v}\left(1-v^2\right)\right.\\ &\left.+r'\,(\kappa\mathbf{e}_{r'} - \mathbf{v})\left(\mathbf{e}_{r'}\cdot\dot{\mathbf{v}}\right) + r'\left(\mathbf{e}_{r'}\left(\mathbf{e}_{r'}\cdot\dot{\mathbf{v}}\right) - \dot{\mathbf{v}}\right)(1-\kappa)\right]\\ = \frac{q}{4\pi\epsilon_0}&\frac{\left(1-v^2\right)(\mathbf{x}' - \mathbf{v}r')}{(r' - \mathbf{x}'\cdot\mathbf{v})^3} + \frac{q}{4\pi\epsilon_0}\frac{(\mathbf{x}' - \mathbf{v}r')\left(\mathbf{x}'\cdot\dot{\mathbf{v}}\right) - \dot{\mathbf{v}}r'\,(r' - \mathbf{x}'\cdot\mathbf{v})}{(r' - \mathbf{x}'\cdot\mathbf{v})^3}.\end{aligned} \tag{12}$$

Taking into account that the numerator in the second term in the latter expression can be rewritten as the double vector product

$$(\mathbf{x}' - \mathbf{v}r') \left(\mathbf{x}' \cdot \dot{\mathbf{v}}\right) - \dot{\mathbf{v}}r' \left(r' - \mathbf{x}' \cdot \mathbf{v}\right) = \left[\mathbf{x}' \times \left[(\mathbf{x}' - r'\mathbf{v}) \times \dot{\mathbf{v}}\right]\right], \tag{13}$$

expression (12) can be recognized (with the identification $q/4\pi\epsilon_0 = e$, $\mathbf{x}' = \mathbf{R}$, $r' = R$) as the expression (63.8) for electric field in Ref. [2]. Thus, expressions in Refs. [2] and [4] are the same.

Acknowledgements

Supported by FAPESP under grants 2013/00840-9, 2013/16592-4 and 2014/08970-1, by RFBR under Project 15-02-00293a, and by the TSU Competitiveness Improvement Program, by a grant from "The Tomsk State University D.I. Mendeleev Foundation Program". A.A.S. thanks also CAPES for support. A.E.S. thanks the University of São Paulo for hospitality extended to him during the period when this work was being performed.

---

[1] R. de Sangro, G. Finocchiaro, P. Patteri, M. Piccollo, and G. Pizzella, Eur. Phys. J. C **75**, 137 (2015).

[2] L.D. Landau, E.M. Lifshitz, *The Classical Theory of Fields.* (GIF-ML, Moscow, 1962; Pergamon Press, Oxford, 1971).

[3] J.D. Jackson, Classical Electrodynamics (Wiley, New York, 1962 – 1975, pp. 654 – 658).

[4] R.P. Feynman, R.B. Leighton and M. Sands, *The Feynman Lectures in Physics* (Addison-Wesley, Reading Massachusetts, 1963), "Electromagnetism II" Ch 21-1. See also http://www.feynmanlectures.caltech.edu/II_21.html#Ch21-S3. .

[5] J.H. Field, Int. J. Mod. Phys. A **23**, 327 (2008); *Retarded electric and magnetic fields of a moving charge: Feynman's derivation of Liénard-Wiechert potentials revisited,* arXiv:0704.1574[physics.class-ph](2007); *Comment on "Measuring propagation speed of Coulomb fields",* arXiv:1506.06630[physics.gen-ph] (2015) (or arXiv:1506.06630v2 [physics.gen-ph] (2016) for completed version) .